



# Determination of the Best Anthropometric Index of Obesity for Prediction of Prehypertension and Hypertension in a Large Population - Based - Study; the Azar- Cohort

Elnaz Faramarzi,<sup>1</sup> Alireza Ostadrahimi,<sup>2</sup> Zeinab Nikniaz,<sup>1</sup> Mohamad Asgari Jafarabadi,<sup>3</sup> Ali Fakhari,<sup>4</sup> and Mohammadhossein Somi<sup>1\*</sup>

<sup>1</sup>Liver and Gastrointestinal Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, IR Iran

<sup>2</sup>Nutrition Research Center, Tabriz University of Medical Sciences, Tabriz, IR Iran

<sup>3</sup>Medical Education Research Center, Department of Statistics and Epidemiology, Faculty of Health, Tabriz University of Medical Sciences, Tabriz, IR Iran

<sup>4</sup>Psychiatry and Behavioral Sciences Research Center, Tabriz University of Medical Sciences, Tabriz, IR Iran

\*Corresponding author: Mohammadhossein Somi, Professor of Gastroenterology, Liver and Gastrointestinal Diseases Research Center, Tabriz University of Medical Sciences, Tabriz, IR Iran. E-mail: mhosseinsina@yahoo.com

Received 2017 August 16; Revised 2017 October 17; Accepted 2017 December 20.

## Abstract

**Background:** Identification of reliable predictors of hypertension and prehypertension in each population is essential for early detection of at-risk people and also planning preventive strategies.

**Objectives:** The present study aimed to assess the abilities of different indices of general and central obesity in the prediction of incident prehypertension and hypertension in a large population-based study in Iran.

**Methods:** In this cross-sectional study, we evaluated the anthropometric and blood pressure measurements in 10,137 subjects (35-70 years) from Azar- Cohort study, north-west of Iran. Blood pressure was measured twice, with 10 minutes apart, from both upper extremities. Height, weight, waist, and hip circumferences measured according to the NIH guidelines. Hypertension considered as systolic blood pressure  $\geq 140$  and diastolic blood pressure  $\geq 90$  mmHg or current use of antihypertensive medication for management of hypertension.

**Results:** The mean BMI ( $\text{kg}/\text{m}^2$ ), WC (cm), waist- to- hip ratio (WHR), and waist- to- height ratio (WHtR) were  $28.75 \pm 4.83$ ,  $94.69 \pm 11.23$ ,  $0.90 \pm 0.07$ , and  $0.58 \pm 0.07$ , respectively. The prevalence of prehypertension and hypertension were 16.3% and 23.3%, respectively. There was a significant association between prehypertension and hypertension with all included anthropometric indices in both men and women (All  $P < 0.001$ ). In men, obesity increased the risk of prehypertension and hypertension by 3.13 (95% CI: 2.48 - 3.94) and 4.06 (95% CI: 3.23 - 5.10), respectively. In women with WHtR  $> 0.5$  cm, the risk of hypertension increased by 6.27 (95% CI: 4.39 - 8.95) times.

**Conclusions:** According to the result of this large population-based study, BMI is the best predictor of prehypertension and hypertension in men and also the waist- to- hip ratio combined with BMI were the best predictors of prehypertension and hypertension in women. These findings may have significant implications in using the most useful screening index for predicting hypertension in Iranian adults and also using waist- to- hip ratio for early detection of pre-hypertension.

**Keywords:** Anthropometric, Cohort Study, Hypertension, Index, Measurements, Obesity, Prehypertension

## 1. Background

Hypertension is the global public-health problem and leading modifiable risk factor for cardiovascular disease (CVD) (1). It is predicted that in 2025, the number of adults with hypertension will increase to 1.56 billion (1). About 2/3 of the patients with hypertension are now living in developing countries (1, 2). Iran, as a Middle Eastern and developing country, faced with rapid population, social, and economic changes that lead to an upward trend in the

prevalence of many CVD risk factors. A previous study has estimated that the prevalence of hypertension in Iranian adults and youth was about 22% (3-5).

Recently, the pre-hypertension rates increase worldwide along with hypertension. In this regard, the findings of Hadaegh et al. indicated that approximately half of Iranians had prehypertension during a ten year follow up (6). Pre-hypertensive subjects are at higher risk of hypertension and CVD compared to normotensive people (7, 8).

Risk factors including harmful use of alcohol, smok-

ing, high dietary salt intake, excess body weight, physical inactivity, and hypercholesterolemia are among the risk factors of hypertension and prehypertension (9-11). Among these risk factors, obesity considered as an independent risk factor for hypertension and CVD. There are a lot of studies that have assessed the relationship between different indices of obesity (such as body mass index (BMI), abdominal obesity, and so on) and hypertension. Although previously it has been shown that the prevalence of hypertension increases with general and central obesity, however, these studies mostly considered BMI and waist circumference (WC) as obesity indices and other predictors of cardiovascular disease were rarely investigated and conflicting results were also reported (12-14). On the other hand, despite the increasing incidence of prehypertension and its association with CVD, scarce studies were conducted about the association between general and abdominal obesity and pre-hypertension. Moreover, the predictive power of different anthropometric indices also depends on the ethnic origin of the studied population. Therefore, defining population-specific anthropometric indices are necessary to determine subjects at risk for hypertension (15-17).

For early detection of at-risk people and also planning of preventive strategies, it is crucial to identify strong predictors of hypertension and prehypertension in each population. Therefore, for the first time in the Iranian population, the power of different indices of general and central obesity in the prediction of incident prehypertension and hypertension were assessed.

## 2. Methods

The data of this cross-sectional study were gained from the Azar-cohort study, which is established in Shabestar in East Azerbaijan province (north-west of Iran) and also it is a part of a large Persian cohort study (The prospective epidemiological research studies of the Iranian Adults). This study was launched in October 2014 and conducted in 3 phases of pilot, enrollment, and follow-up. Written informed consent was obtained from all participants. This prospective study was approved by Ethics Committee of Tabriz University of Medical Sciences (tbzmed.rec.1393.205), Tabriz, Iran.

### 2.1. Subjects and Data

All eligible individuals 35 - 70 years of age in the region of Shabestar are invited to participate in the study, and those who signed the consent form are included (totally 10,137 subjects). Those included were inhabitants in Shabestar for at least nine months. The participants with severe psychiatric or physical illness were excluded from

the study. From October 2014 to June 2016, out of 10,788 subjects who were invited, 10,137 responded. The response rate was 94%.

In the Azar cohort study, firstly the lists of all subjects age 35 - 70 years were obtained from the health centers of Shabestar region. From small cities and villages, approximately all eligible individuals 35 - 70 years were invited. In other cities, more than 60% of eligible subjects were invited. In January 2017, the enrollment phase has been finished and 15,000 subjects were recruited. For invitation, 1 week before the day of visit, people were contacted by telephone and conditions of the cohort study were explained. In addition, if the person agreed to participate in the study, the date of visit day, were given to him/her.

### 2.2. Anthropometric Measurements

Mounted tape was used for measuring the height to the nearest 1 mm and Seca scale was used for recording weight to the nearest 0.1 kg, according to standard protocols. Weight was measured by Seca scale. Every day, suitable standard 10 kg weights was used for calibration of weighing scales. Body mass index was calculated by dividing weight (in kilogram) by the square of height (in meter). BMI was classified according to the WHO (18) categories: underweight (BMI < 18.5 kg/m<sup>2</sup>), normal (BMI 18.5 - 24.9 kg/m<sup>2</sup>), overweight (BMI 25 - 29.9 kg/m<sup>2</sup>), and obesity (BMI ≥ 30 kg/m<sup>2</sup>). WC and hip circumference of subjects are measured according to NIH (National Institute of Health) guidelines. Waist to hip ratio (WHR) and waist to height ratio (WHtR) were calculated by dividing waist to hip and height, respectively. In men, WC was ≥ 102 cm and WHR ≥ 0.9 cm, in women WC was ≥ 88 cm and WHR ≥ 0.85 cm, and in both genders, WHtR ≥ 0.5 were considered as a risk factor.

### 2.3. Blood Pressure Definition

Blood pressure was measured twice with 10 minutes apart from both arms separately, by a trained nurse using a mercury sphygmomanometer (Riester, Germany). It was calibrated by mercury sphygmomanometers.

The mean of measurements was recorded as the blood pressure. Prehypertension was defined as having either a Systolic Blood Pressure of 120 to 139 mmHg and/or Diastolic Blood Pressure of 80 to 89 mmHg, according to JNC7, in persons who were not on antihypertensive medication (19). Subjects with SBP ≥ 140 and DBP ≥ 90 mmHg or use of an antihypertensive drug with a history of hypertension were considered as patients with hypertension.

To reduce individual errors in anthropometric and blood pressure measurements sections, in each section, only one trained independent person assessed anthropometric and blood pressure measurements. We have

an epidemiologist who checked all filled questionnaires as well as anthropometric and blood pressure measurements. Moreover, another one monitored the work of the epidemiologist.

#### 2.4. Statistical Analysis

Statistical package for the social sciences (SPSS Inc. version 16.0, Chicago, IL, USA) was used to analyze data. Descriptive statistics were obtained for all study variables and reported as mean  $\pm$  SD and also number (percent) where applicable. The normality of data was assessed by the Kolmogorov-Smirnov test. All data had normal distribution.

We used logistic regression analysis for estimating crude and adjusted odds ratios (OR) and their corresponding 95% confidence intervals (95%CI). On the basis of BMI classification, 13 women and 47 men were underweight that were excluded in logistic regression analysis. A P value of less than 0.05 was considered statistically significant.

### 3. Results

The baseline characteristics of study subjects stratified by sex were presented in Table 1. The mean age of the participants was  $49.32 \pm 9.15$  years. The mean BMI ( $\text{kg}/\text{m}^2$ ), WC (cm), WHR, and WHtR were  $28.75 \pm 4.83$ ,  $94.69 \pm 11.23$ ,  $0.90 \pm 0.07$ , and  $0.58 \pm 0.07$ , respectively. About 36.6% of participants were obese, and 23.3% of them had hypertension.

Table 2 presents the percent of patients with different blood pressure status in different anthropometric subclasses. As can be seen, respectively 32.5%, 32.6%, 28.6%, and 25.6% of subjects with obesity,  $\text{WC} \geq 102$  cm in men and  $>88$  cm in women,  $\text{WHR} \geq 0.9$  (men) and 0.85 (women) and  $\text{WHtR} > 0.5$ , had hypertension. The prevalence of hypertension in obese women, women with  $\text{WC} \geq 88$  cm,  $\text{WHR} \geq 0.85$ , and  $\text{WHtR} \geq 0.5$  were more than their men counterparts.

The association between different anthropometric indices and prehypertension and hypertension was presented in Table 3. There is a significant association between prehypertension and hypertension with all included anthropometric indices in both genders (All  $P < 0.001$ ).

In men, in an unadjusted model, BMI had the highest OR with prehypertension and WHtR had the highest OR with hypertension. However, after adjusting for age, BMI had the highest association with both hypertension and prehypertension. Moreover, obesity increased the risk of prehypertension and hypertension by 3.13 (95%CI: 2.48 - 3.94) and 4.06 (95%CI: 3.23 - 5.10). After adjustment for age, the observed association were increased (OR = 3.43; 95% CI: 2.70 - 4.35, OR = 4.74; 95%, CI = 3.71 - 6.07). After WHtR, BMI,

Table 1. Baseline Characteristics of Participants

Variable	Men	Women	Total
Age, y	49.92 $\pm$ 9.12	48.83 $\pm$ 9.14	49.32 $\pm$ 9.15
Weight, kg	79.58 $\pm$ 13.50	72.98 $\pm$ 12.83	75.95 $\pm$ 13.54
Height, cm	170.51 $\pm$ 6.72	172.98 $\pm$ 12.83	162.63 $\pm$ 9.70
Waist, cm	96.09 $\pm$ 11.22	93.55 $\pm$ 11.11	94.69 $\pm$ 11.23
BMI, $\text{kg}/\text{m}^2$	27.35 $\pm$ 4.27	29.90 $\pm$ 4.96	28.75 $\pm$ 4.83
WHR	0.93 $\pm$ 0.07	0.87 $\pm$ 0.07	0.90 $\pm$ 0.07
WHtR	0.56 $\pm$ 0.06	0.59 $\pm$ 0.07	0.58 $\pm$ 0.07
SBP, mm/Hg	112.34 $\pm$ 16.59	113.85 $\pm$ 17.62	113.17 $\pm$ 17.18
DBP, mm/Hg	73.22 $\pm$ 9.32	73.29 $\pm$ 10.07	73.26 $\pm$ 9.74
BMI <sup>a</sup> , $\text{kg}/\text{m}^2$			
Underweight	48 (1.1)	13 (0.2)	61 (0.6)
Normal (18.5 - 24.9)	1292 (28.4)	835 (15)	2127 (21)
Overweight (25 - 29.9)	2081 (45.78)	2152 (38.5)	4233 (41.8)
Obese, ( $\geq 30$ )	1127 (24.7)	2576 (46.1)	3703 (36.6)
WC <sup>a</sup> , cm			
< 102 in men/ < 88 in women	3275 (72)	1734 (31.1)	5009 (49.5)
$\geq 102$ in men/ $> 88$ in women	127 (28)	3834 (68.8)	515 (50.5)
WHR <sup>a</sup>			
< 0.9 in men/ < 0.85 in women	1357 (29.8)	2033 (36.4)	3390 (33.4)
$\geq 0.9$ in men/ $\geq 0.85$ in women	3190 (70)	3542 (63.4)	6732 (66.4)
WHtR <sup>a</sup>			
< 0.5	774 (17)	469 (8.4)	1243 (12.3)
$\geq 0.5$	3773 (82.9)	5107 (91.5)	8880 (87.6)
Hypertension systolic/diastolic <sup>a</sup> , mmHg			
Normal (< 120 and < 80)	2968 (65.2)	3117 (55.8)	6085 (60)
Prehypertension (120 - 139 or 80 - 89)	731 (16.1)	919 (16.5)	1650 (16.3)
Hypertension, $\geq 140$ or $\geq 90$ )	831 (18.2)	1530 (27.4)	2361 (23.3)

Abbreviations: BMI, body mass index; DPB, diastolic blood pressure; N, number; SBP, systolic blood pressure; WC, waist circumference; WHR, waist to hip ratio; WHtR, waist to height ratio.

<sup>a</sup>Values are expressed as No. (%).

WHR, and WC, respectively had the highest OR for hypertension.

In women, BMI and WHtR had the highest associations with prehypertension and hypertension in both adjusted

**Table 2.** The Percent of Patients with Different Blood Pressure Status in Different Anthropometric Subclass<sup>a</sup>

Variables	Men			Women			Total		
	Normal	Prehypertension (120 - 139 or 80 - 89)	Hypertension ( $\geq 140$ or $\geq 90$ )	Normal (< 120 and < 80)	Prehypertension (120 - 139 or 80 - 89)	Hypertension ( $\geq 140$ or $\geq 90$ )	Normal (< 120 and < 80)	Prehypertension (120 - 139 or 80 - 89)	Hypertension ( $\geq 140$ or $\geq 90$ )
<b>Body mass index, kg/m<sup>2</sup></b>									
Normal (18.5 - 24.9)	1008 (78.1)	139 (10.8)	133 (10.3)	611 (73.3)	90 (10.8)	130 (15.6)	1619 (76.2)	229 (10.8)	263 (12.4)
Over-weight (25 - 29.9)	1351 (64.2)	338 (16.2)	385 (18.5)	1330 (61.8)	318 (14.8)	501 (23.3)	2681 (63.3)	656 (15.5)	886 (20.9)
Obese ( $\geq 30$ )	572 (50.7)	247 (21.9)	307 (27.2)	1159 (45)	510 (19.8)	898 (34.8)	1731 (46.7)	757 (20.4)	1205 (32.5)
<b>Waist circumference (WC)</b>									
< 102 in men/ < 88 in women	2342 (71.5)	496 (14.3)	448 (13.7)	1289 (74.3)	196 (11.3)	245 (14.1)	363 (72.5)	665 (13.3)	693 (13.8)
$\geq 102$ in men/ $\geq 88$ in women	625 (49.1)	261 (20.5)	383 (30.11)	1827 (47.5)	722 (18.8)	12.84 (33.4)	2452 (47.9)	983 (19.2)	1667 (32.6)
<b>Waist to hip ratio (WHR)</b>									
< 0.9 in men and < 0.85 in women	1077 (79.3)	152 (11.2)	120 (8.8)	1465 (72.1)	245 (12.1)	313 (15.4)	2542 (75)	397 (11.7)	433 (12.8)
$\geq 0.9$ in men and $\geq 0.85$ in women	1890 (59.2)	578 (18.1)	711 (22.3)	1216 (46.6)	673 (19.4)	1216 (34.3)	3540 (52.6)	1251 (18.6)	1927 (28.6)
<b>Waist-to-height ratio (WHtR)</b>									
< 0.5	649 (83.9)	65 (8.41)	53 (6.8)	389 (82.9)	44 (9.4)	34 (7.2)	1038 (83.5)	109 (8.8)	87 (7)
$\geq 0.5$	2318 (61.4)	665 (17.6)	778 (20.6)	2726 (53.4)	874 (17.1)	1495 (29.3)	5044 (56.8)	1539 (17.3)	2273 (25.6)

<sup>a</sup> Normal: systolic blood pressure < 120 mmHg and diastolic blood pressure < 80 mmHg; Prehypertension: systolic blood pressure < 120 - 139 mmHg and diastolic blood pressure < 80 - 89 mmHg; hypertension, systolic blood pressure  $\geq 140$  mmHg and diastolic blood pressure  $\geq 90$  mmHg.

and unadjusted models. In women with WHtR > 0.5 cm, the risk of hypertension increased by 6.27 (95%CI: 4.39 - 8.95) times. However, this association was decreased after adjustment for age.

After adjustment for age, the strong predictor of prehypertension and hypertension were BMI and WHtR in men and women, respectively.

#### 4. Discussion

This paper presents the cross-sectional analysis of the 1st phase of a comprehensive community-based survey of Azar cohort conducted in North West Iran. According to the results, in the large sample of Azar cohort, the prevalence of prehypertension and hypertension were 16.3% and 23.3%, respectively. The prevalence of hypertension in East Azerbaijan was reported to be 20.47% in 2007 (20), which indicates that prehypertension/hypertension rates in adults are rising at an alarming rate. Previous studies in the USA (21), China (22), and Egypt (23) reported a

higher prevalence of prehypertension and hypertension. Moreover, in the present study, the prevalence of hypertension in subjects with BMI  $\geq 30$  kg/m<sup>2</sup>, WC  $\geq 102$  cm in men and  $\geq 88$  cm in women, WHR  $\geq 0.9$  (men) and 0.85 (women), and WHtR  $\geq 0.5$  were 32.5%, 32.6%, 28.6%, and 25.6%, respectively, which was lower than previous reports from other countries. In the Indian population, the prevalence of hypertension was reported to be 40.8% (BMI > 27.6 kg/m<sup>2</sup>), 47.2% (WC > 98.1 cm), 44.7% (WHR > 0.98), and 45.9% (WHtR > 0.59) (24). The prevalence of prehypertension in Vietnamese adults was reported to be 40.7% (men) and 38.2% (women), and the prevalence of hypertension was 50% (men) and 34.9% (women). A wide range of prevalence of hypertension in various countries may be due to the differences in the social and cultural differences, different behavioral and dietary lifestyles, the age span, as well as the methodology used.

The results of the present study further support the idea of an association between anthropometric indices and hypertension. In a large study of the Chinese pop-

**Table 3.** Logistic Regression of the Association Between Prehypertension and Hypertension and Anthropometric Indices in Iranian Adults<sup>a</sup>

Variables	Pre-Hypertension		Hypertension	
	Unadjusted OR <sup>b</sup>	Adjusted OR <sup>c</sup>	Unadjusted OR <sup>b</sup>	Adjusted OR <sup>c</sup>
<b>Men</b>				
<b>BMI, kg/m<sup>2</sup></b>				
18.5 - 24.99	1	1	1	1
25 - 29.99	1.81 (1.46, 2.24)	1.90 (1.52, 2.36)	2.16 (1.74, 2.67)	2.54 (1.88, 3.42)
≥ 30	3.13 (2.48, 3.94)	3.43 (2.70, 4.35)	4.06 (3.23, 5.10)	4.74 (3.71, 6.07)
<b>WC, cm</b>				
< 102	1	1	1	1
≥ 102	2.08 (1.75, 2.48)	2.01 (1.68, 2.40)	3.20 (2.72, 3.76)	2.93 (2.46, 3.49)
<b>WHR</b>				
< 0.9	1	1	1	1
≥ 0.9	2.16 (1.78, 2.63)	1.94 (1.59, 2.36)	3.37 (2.74, 4.15)	2.76 (2.22, 3.43)
<b>WHtR</b>				
< 0.5	1	1	1	1
≥ 0.5	2.86 (2.18, 3.74)	2.64 (2.01, 3.47)	4.11 (3.07, 5.50)	3.63 (2.68, 4.92)
<b>Women</b>				
<b>BMI, kg/m<sup>2</sup></b>				
18.5 - 24.99	1	1	1	1
25 - 29.99	1.62 (1.26, 2.09)	1.61 (1.25, 2.09)	1.77 (1.42, 2.19)	1.86 (1.46, 2.37)
≥ 30	2.98 (2.33, 3.81)	2.49 (2.01, 3.07)	3.44 (2.95, 4.48)	3.74 (2.96, 4.74)
<b>WC, cm</b>				
< 88	1	1	1	1
≥ 88	2.59 (2.18, 3.09)	2.32 (1.95, 2.77)	3.69 (3.16, 4.31)	2.87 (2.43, 3.41)
<b>WHR</b>				
< 0.85	1	1	1	1
≥ 0.85	2.43 (2.07, 2.86)	2.00 (1.69, 2.37)	3.44 (2.99, 3.97)	2.09 (1.79, 2.45)
<b>WHtR</b>				
< 0.5	1	1	1	1
≥ 0.5	2.83 (2.05, 3.90)	2.36 (1.70, 3.26)	6.27 (4.39, 8.95)	4.26 (2.91, 6.22)

Abbreviations: OR, odds ratio; WC, waist circumference; WHR, waist to hip ratio; WHtR, waist to height ratio.

<sup>a</sup>Prehypertension: systolic blood pressure < 120 - 139 mmHg and diastolic blood pressure < 80 - 89 mmHg; hypertension: systolic blood pressure ≥ 140 mmHg and diastolic blood pressure ≥ 90 mmHg.

<sup>b</sup>95%CI: confidence interval.

<sup>c</sup>Adjusted for age.

ulation (25), BMI and WC were associated with hypertension. In the mentioned study, other anthropometric indices such as WHtR and WHR as well as the association between prehypertension and anthropometric indices were not studied. In a church-based study on the Nigerian population, BMI and WHtR were significantly associated with hypertension (26). In addition, they found that BMI and WC had more predictor power for prehypertension (26).

Kaur et al. in industrial men showed that both hyper-

tension and prehypertension were significantly associated with BMI, WC, WHR, and WHtR (24). Multiple mechanisms, including the renin-angiotensin-aldosterone system activation, increased sympathetic nervous system (SNS) activity and insulin resistance, increased renal sodium reabsorption, impaired pressure natriuresis, and volume expansion were involved in obesity-related HTN association (27).

Although the association between obesity, prehyper-

tension, and hypertension were reported in previous studies from Iran and other countries, the power of each anthropometric indices in predicting prehypertension and hypertension should be determined in each population. As far we know, no study on the Iranian population has been performed to answer the question, “which of the anthropometric indices were the best predictor of prehypertension and hypertension”. The results of the present study showed that WHtR is correlated with a higher OR than BMI for hypertension, in contrast, BMI is associated with a higher OR for prehypertension. In line with our study, Deng et al. in a study on the Chinese population showed that BMI had a higher association with prehypertension, while WC had higher an OR than BMI for hypertension (25). The strong correlation was observed between BMI and prehypertension and hypertension in the Chinese population (28). Moreover, according to the results of the DECODA study, both waist-to-height ratios and BMI were reported as a strong predictor of hypertension in Asians (29). In contrast, Ma et al. reported that WC was more predictive than BMI and WHtR for prehypertension (30). In Australian adults, WHR has been suggested as the most useful index of obesity to recognize individuals with cardiovascular disease risk factors (24). The discrepancies between the different studies results may be ascribed to the differences in the study populations. It seems that WHtR along with BMI is a useful index for recognizing prehypertension and hypertension, rather than using only one of these measures for the Iranian population. WHtR and BMI are the convenient index for use in the population. They have only one cut-off for both sexes, which is easy to memorize and consumer-friendly.

As reported by previous studies, age had modifying effects on the association between different anthropometric indices and hypertension (25, 31). These observations may be due to increasing the central obesity and declining in height with an increase in age.

The strengths of the present study were using a large population-based sample and prospective cohort design, duplicate measuring of anthropometric indices, and blood pressure. However, considering insufficient covariates included in the present study, these results should be interpreted with caution. Other covariates, rather than age, may influence the association between anthropometric indices and hypertension such as depression, anxiety, dietary factors, and physical activity pattern. Moreover, due to the cross-sectional analysis of data, causal relations may not be established.

In conclusion, the results of this large population-based study showed that all anthropometric indices were associated with prehypertension and hypertension both in men and women. Moreover, BMI is the best predictor of

prehypertension and hypertension in men. Besides, WHtR combined with BMI was the best predictor of prehypertension and hypertension in women. These findings may have significant implications in using the most useful screening index for predicting hypertension in Iranian adults and also using WHtR for early detection of pre-hypertension (before hypertension will exist).

## Acknowledgments

The authors are grateful for the financial support of the liver and gastrointestinal disease research center, Tabriz University of Medical Sciences. The authors also are deeply indebted to all subjects who participated in this study. We appreciate the contribution by the investigators and the staff of the Azar cohort study. We thank the close collaboration of the Shabestar health center. Also, we would like to thank the Persian cohort study staff for their technical support.

## Footnotes

**Authors' Contribution:** Study concept and design: Mohammadhossein Somi, Elnaz Faramarzi; acquisition of data: Mohammadhossein Somi; analysis and interpretation of data: Elnaz Faramarzi; drafting of the manuscript: Elnaz Faramarzi, Zeinab Nikniaz; critical revision of the manuscript for important intellectual content: Ali Fakhari, Zeinab Nikniaz; statistical analysis: Mohamad Asgari Jafarabadi; administrative, technical, and material support: Alireza Ostadrahimi; study supervision: Alireza Ostadrahimi.

**Conflict of Interests:** The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**Funding/Support:** Liver and gastrointestinal diseases research center, Tabriz University of Medical Sciences, Tabriz, Iran.

## References

1. Murray CJL, Lopez AD. *Global health statistics: Global burden of disease and injury series*. Boston: Harvard school of public health; 1996.
2. WHO; Consultation FAO Expert. *Diet, nutrition and the prevention of chronic diseases*. 2003.
3. Abdollahi AA, Bazrafshan HR, Salehi A, Behnampour N, Hoseini SA, Rahmanipour AH, et al. Epidemiology of hypertension among urban population in Golestan province in north of Iran. *J Gorgan Uni Med Sci*. 2007;8:37-41.
4. Janghorbani M, Amini M, Gouya MM, Delavari A, Alikhani S, Mahdavi A. Nationwide survey of prevalence and risk factors of prehypertension and hypertension in Iranian adults. *J Hypertens*. 2008;26(3):419-26. doi:10.1097/HJH.0b013e3282f2d34d. [PubMed: 18300850].

5. Tabrizi JS, Sadeghi-Bazargani H, Farahbakhsh M, Nikniaz L, Nikniaz Z. Prevalence and Associated Factors of Prehypertension and Hypertension in Iranian Population: The Lifestyle Promotion Project (LPP). *PLoS One*. 2016;**11**(10). e0165264. doi: [10.1371/journal.pone.0165264](https://doi.org/10.1371/journal.pone.0165264). [PubMed: [27783691](https://pubmed.ncbi.nlm.nih.gov/27783691/)].
6. Hadaegh F, Hashemina M, Abdi H, Khalili D, Bozorgmanesh M, Arshi B, et al. Prehypertension Tsunami: A Decade Follow-Up of an Iranian Adult Population. *PLoS One*. 2015;**10**(10). e0139412. doi: [10.1371/journal.pone.0139412](https://doi.org/10.1371/journal.pone.0139412). [PubMed: [26439847](https://pubmed.ncbi.nlm.nih.gov/26439847/)].
7. Li G, Guo G, Wang W, Wang K, Wang H, Dong F, et al. Association of prehypertension and cardiovascular risk factor clustering in Inner Mongolia: a cross-sectional study. *BMJ*. 2017;**7**(6). e15340. doi: [10.1136/bmjopen-2016-015340](https://doi.org/10.1136/bmjopen-2016-015340).
8. Tajeu GS, Booth JN, Colantonio LD, Gottesman RF, Howard G, Lackland DT, et al. Incident cardiovascular diseases among adult with blood pressure <140/90 mmHg. *Circulation*. 2017;**136**(9):798–812. doi: [10.1016/S0140-6736\(08\)60655-8](https://doi.org/10.1016/S0140-6736(08)60655-8). [PubMed: [18456100](https://pubmed.ncbi.nlm.nih.gov/18456100/)].
9. World Health Organization. *Global action plan for the prevention and control of noncommunicable diseases 2013-2020*. Geneva: World Health Organization; 2015.
10. Murtaugh MA, Beasley JM, Appel LJ, Guenther PM, McFadden M, Greene T, et al. Relationship of sodium intake and blood pressure varies with energy intake: secondary analysis of the DASH (dietary approaches to stop hypertension)-sodium trial. *Hypertension*. 2018;**71**(5):858–65. doi: [10.1161/HYPERTENSIONAHA.117.10602](https://doi.org/10.1161/HYPERTENSIONAHA.117.10602). [PubMed: [29555665](https://pubmed.ncbi.nlm.nih.gov/29555665/)].
11. Stenehjem JS, Hjerkind KV, Nilsen TIL, Amiri B, Majzoobi M, Erfani H, et al. Diposity, physical activity and risk of hypertension :prospective data from the population -based HUNT study, Norway. *J Hum Hypertens*. 2018;**32**(4):278–86. doi: [10.1038/s41371-018-0042-5](https://doi.org/10.1038/s41371-018-0042-5). [PubMed: [29483587](https://pubmed.ncbi.nlm.nih.gov/29483587/)].
12. Browning LM, Hsieh SD, Ashwell M. A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 0.5 could be a suitable global boundary value. *Nutr Res Rev*. 2010;**23**(2):247–69. doi: [10.1017/S0954422410000144](https://doi.org/10.1017/S0954422410000144). [PubMed: [20819243](https://pubmed.ncbi.nlm.nih.gov/20819243/)].
13. Nurdiantami Y, Watanabe K, Tanaka E, Pradono J, Anme T. Association of general and central obesity with hypertension. *Clin Nutr*. 2018;**37**(4):1259–63. doi: [10.1016/j.clnu.2017.05.012](https://doi.org/10.1016/j.clnu.2017.05.012). [PubMed: [28583324](https://pubmed.ncbi.nlm.nih.gov/28583324/)].
14. Tran NTT, Blizzard CL, Luong KN, Truong NLV, Tran BQ, Otahal P, et al. The importance of waist circumference and body mass index in cross-sectional relationships with risk of cardiovascular disease in Vietnam. *Plos one*. 2018;**13**(5):908–16. doi: [10.1371/journal.pone.0198202](https://doi.org/10.1371/journal.pone.0198202). [PubMed: [29813112](https://pubmed.ncbi.nlm.nih.gov/29813112/)].
15. Esmailzadeh A, Mirmiran P, Azizi F. Waist-to-hip ratio is a better screening measure for cardiovascular risk factors than other anthropometric indicators in Tehranian adult men. *Int J Obes Relat Metab Disord*. 2004;**28**(10):1325–32. doi: [10.1038/sj.ijo.0802757](https://doi.org/10.1038/sj.ijo.0802757). [PubMed: [15314626](https://pubmed.ncbi.nlm.nih.gov/15314626/)].
16. Harris MM, Stevens J, Thomas N, Schreiner P, Folsom AR. Associations of fat distribution and obesity with hypertension in a bi-ethnic population: the ARIC study. Atherosclerosis Risk in Communities Study. *Obes Res*. 2000;**8**(7):516–24. doi: [10.1038/oby.2000.64](https://doi.org/10.1038/oby.2000.64). [PubMed: [11068957](https://pubmed.ncbi.nlm.nih.gov/11068957/)].
17. Wang Z, Hoy WE. Body size measurements as predictors of type 2 diabetes in Aboriginal people. *Int J Obes Relat Metab Disord*. 2004;**28**(12):1580–4. doi: [10.1038/sj.ijo.0802771](https://doi.org/10.1038/sj.ijo.0802771). [PubMed: [15356663](https://pubmed.ncbi.nlm.nih.gov/15356663/)].
18. World Health Organization. *BMI Classification*. 2011.
19. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JJ, et al. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension*. 2003;**42**(6):1206–52. doi: [10.1161/01.HYP.0000107251.49515.c2](https://doi.org/10.1161/01.HYP.0000107251.49515.c2). [PubMed: [14656957](https://pubmed.ncbi.nlm.nih.gov/14656957/)].
20. Ahmadi A, Mobasheri M, Soori H. Prevalence of major coronary heart disease risk factors in Iran. *Int J Epidemiol Res*. 2014;**1**(1):3–8.
21. Gupta AK, McGlone M, Greenway FL, Johnson WD. Prehypertension in disease-free adults: a marker for an adverse cardiometabolic risk profile. *Hypertens Res*. 2010;**33**(9):905–10. doi: [10.1038/hr.2010.91](https://doi.org/10.1038/hr.2010.91). [PubMed: [20535113](https://pubmed.ncbi.nlm.nih.gov/20535113/)].
22. Zhang M, Batu B, Tong W, Li H, Lin Z, Li Y, et al. Prehypertension and cardiovascular risk factor clustering among Mongolian population in rural and animal husbandry area, Inner Mongolia, China. *Circ J*. 2009;**73**(8):1437–41. [PubMed: [19564702](https://pubmed.ncbi.nlm.nih.gov/19564702/)].
23. Gulens M. Egyptians score high on blood pressure. *BMJ*. 1994;**308**(6925):360. doi: [10.1136/bmj.308.6925.360](https://doi.org/10.1136/bmj.308.6925.360).
24. Kaur P, Radhakrishnan E, Sankarasubbaiyan S, Rao SR, Kondalsamy-Chennakesavan S, Rao TV, et al. A comparison of anthropometric indices for predicting hypertension and type 2 diabetes in a male industrial population of Chennai, South India. *Ethnic Dis*. 2008;**18**(1):31–6.
25. Deng WW, Wang J, Liu MM, Wang D, Zhao Y, Liu YQ, et al. Body mass index compared with abdominal obesity indicators in relation to prehypertension and hypertension in adults: the CHPSNE study. *Am J Hypertens*. 2013;**26**(1):58–67. doi: [10.1093/ajh/hps001](https://doi.org/10.1093/ajh/hps001). [PubMed: [23382328](https://pubmed.ncbi.nlm.nih.gov/23382328/)].
26. Ononamadu CJ, Ezekwesili CN, Onyeukwu OF, Umeogaju UF, Ezeigwe OC, Ihegboro GO. Comparative analysis of anthropometric indices of obesity as correlates and potential predictors of risk for hypertension and prehypertension in a population in Nigeria. *Cardiovasc J Afr*. 2017;**28**(2):92–9. doi: [10.5830/CVJA-2016-061](https://doi.org/10.5830/CVJA-2016-061). [PubMed: [27701484](https://pubmed.ncbi.nlm.nih.gov/27701484/)].
27. Kurukulasuriya LR, Stas S, Lastra G, Manrique C, Sowers JR. Hypertension in obesity. *Med Clin North Am*. 2011;**95**(5):903–17. doi: [10.1016/j.mcna.2011.06.004](https://doi.org/10.1016/j.mcna.2011.06.004). [PubMed: [21855699](https://pubmed.ncbi.nlm.nih.gov/21855699/)].
28. Hu L, Huang X, You C, Li J, Hong K, Li P, et al. Prevalence and Risk Factors of Prehypertension and Hypertension in Southern China. *PLoS One*. 2017;**12**(1). e0170238. doi: [10.1371/journal.pone.0170238](https://doi.org/10.1371/journal.pone.0170238). [PubMed: [28095471](https://pubmed.ncbi.nlm.nih.gov/28095471/)].
29. Decoda Study G, Nyamdorj R, Qiao Q, Lam TH, Tuomilehto J, Ho SY, et al. BMI compared with central obesity indicators in relation to diabetes and hypertension in Asians. *Obesity (Silver Spring)*. 2008;**16**(7):1622–35. doi: [10.1038/oby.2008.73](https://doi.org/10.1038/oby.2008.73). [PubMed: [18421260](https://pubmed.ncbi.nlm.nih.gov/18421260/)].
30. Ma M, Tan X, Zhu S. Prehypertension and its optimal indicator among adults in Hubei Province, Central China, 2013–2015. *Clin Exp Hypertension*. 2017;**39**(6):532–8.
31. Benetou V, Bamia C, Trichopoulos D, Mountokalakis T, Psaltopoulou T, Trichopoulou A. The association of body mass index and waist circumference with blood pressure depends on age and gender: a study of 10,928 non-smoking adults in the Greek EPIC cohort. *Eur J Epidemiol*. 2004;**19**(8):803–9. [PubMed: [15469038](https://pubmed.ncbi.nlm.nih.gov/15469038/)].