



# The Effect of Rutin on Progesterone and Estrogen Receptor Expression in Uterine Endometrial Tissue in the Heterotopic Transplantation of Newborn Mouse Ovary

Tayebeh Hadigol<sup>1</sup>, Aligholi Sobhani<sup>2</sup>, Masoud Hemadi<sup>1</sup>, Saeid Nekoonam<sup>2</sup>, Alireza Shams<sup>3</sup>, Bahram Eslami Farsani<sup>1</sup>, Maryam Dastoorpoor<sup>4</sup> and Ghasem Saki<sup>1,\*</sup>

<sup>1</sup>Department of Anatomy, Faculty of Medicine, Jundishapur University of Medical Sciences, Ahvaz, Iran

<sup>2</sup>Department of Anatomy, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

<sup>3</sup>Department of Anatomy, Faculty of Medicine, Alborz University of Medical Sciences, Karaj, Iran

<sup>4</sup>Department of Epidemiology and Biostatistics, Menopause Andropause Research Center, Jundishapur University of Medical Sciences, Ahvaz, Iran

\*Corresponding author: Department of Anatomy, Cellular and Molecular Research Center, Faculty of Medicine, Jundishapur University of Medical Sciences, Ahvaz, Iran. Tel: +98-6133738628, E-mail: ghasemsaki@yahoo.com

Received 2018 November 11; Revised 2019 April 18; Accepted 2019 April 21.

## Abstract

**Background:** Rutin (quercetin-3-rhamnosyl-glucoside), a flavonoid, is derived from plants and has antioxidant properties.

**Objectives:** This study aimed to evaluate the effect of different concentrations of rutin on mouse ovary heterotopic allotransplantation.

**Methods:** The present animal experimental study was conducted on 40 female adult Balb/c mice weighing  $30 \pm 5$  g at the Jundishapur University of Medical Sciences, Ahvaz, Iran, during 2016 - 2018. The mice were divided by permuted block randomization into 8 groups ( $n = 5$ ): OVX (ovariectomy), as the negative control; normal (positive control); OVX + OVA (ovariectomy and transplantation) (control), treated with 0.5 mL of normal saline; OVX + OVA + 10 mg/kg of rutin; OVX + OVA + 30 mg/kg of rutin; OVX + OVA + 60 mg/kg of rutin; OVX + OVA + 100 mg/kg of rutin; and the autograft. Groups were treated daily. Fourteen days after transplantation, ovarian grafts were collected and processed histologically for follicle number counting. Serum estrogen ( $E_2$ ) and progesterone ( $P_4$ ) levels were evaluated. Furthermore, the expression of Estrogen Receptor alpha ( $ER\alpha$ ), Estrogen Receptor beta ( $ER\beta$ ), and Progesterone Receptor (PR) in the uterine endometrial tissue was tested using qRT-PCR and western blotting.

**Results:** A decrease in the number of mature follicles and increase in the number of atretic follicles (mean  $\pm$  SD: OVX + OVA + 30 =  $19.00 \pm 1.000$ , OVX + OVA + 60 =  $25.00 \pm 5.000$ , and OVX + OVA + 100 =  $23.00 \pm 2.646$ ) were observed in all groups treated with rutin in comparison with the control group (mean  $\pm$  SD:  $12.33 \pm 2.517$ ) ( $P$  value < 0.05). The level of  $E_2$  and  $P_4$  (mean  $\pm$  SD: OVX + OVA + 100 =  $6.133 \pm 1.026$ ) increased in comparison with the OVX + OVA group (mean  $\pm$  SD:  $0.4667 \pm 0.2517$ ) ( $P$  value < 0.05). The protein expression of  $ER\alpha$  (mean  $\pm$  SD: OVX + OVA + 10 =  $1.615 \pm 0.1701$  and OVX + OVA + 30 =  $1.744 \pm 0.1779$ ) in comparison with the control group (mean  $\pm$  SD:  $0.7089 \pm 0.1131$ ), and  $ER\beta$  (mean  $\pm$  SD: OVX + OVA + 10 =  $0.7747 \pm 0.4365$ , OVX + OVA + 30 =  $0.9220 \pm 0.1245$ , OVX + OVA + 60 =  $0.7701 \pm 0.2150$ , and OVX + OVA + 100 =  $0.6676 \pm 0.1547$ ) increased in a dose-dependent manner in all groups treated with rutin in comparison with the OVX + OVA group (mean  $\pm$  SD:  $0.1534 \pm 0.06109$ ) ( $P$  value < 0.05). No significant changes in PR were found in groups treated with rutin in comparison with the control group.

**Conclusions:** The results of the present study indicated that rutin increases  $E_2$  and  $P_4$  levels in ovarian hetero allograft mice. Rutin also upregulated the expression of  $ER\alpha$  and  $ER\beta$  but had no significant effect on PR.

**Keywords:** Allografts, Estrogen Receptor Alpha, Estrogen Receptor Beta, Follicle, Mice, Ovary, Polymerase Chain Reaction, Progesterone, Rutin, Transplantation, Up-Regulation, Western Blot

## 1. Background

The risk of sterility is very serious in young women with cancer. Upon the progress in treatments, there are now several options available for fertility preservation, and thus the use of transplantation in reproductive medicine has been considered. Despite being an attractive and help-

ful method in reproductive medicine in the early days, there are some problems with ovarian transplantation (1). The most important challenge in ovarian tissue transplantation is the survival rate and satisfaction of the functional longevity of the transplanted ovary (2).

During the engraftment and neovascularization of the

transplanted ovarian tissue, an initial ischemia usually occurs, followed by a reperfusion period, which leads to the production of reactive oxygen species (ROS) with oxidative properties (3, 4). Superoxide and hydroxyl radicals are involved in tissue injury through the initiation of lipid peroxidation and damage the cell membrane in the transplanted tissue (5). ROS products are also directly responsible for the oxidative damage of cellular structures, such as DNA, RNA, proteins, and lipid ingredients in ischemic tissues (6). The mentioned events can lead to follicular loss and engraftment insufficiency in the transplanted ovarian tissue (7-9). Due to the critical effect of oxidative stress in transplantation, as described above, antioxidants can be applied to reduce the production of free radicals. Such drugs with antioxidant properties are able to counter the consequences of ischemia and reperfusion injury after ovarian tissue transplantation.

Rutin (quercetin-3-rhamnosyl-glucoside) is a flavonol, abundantly found in apples, tea, onions, and other plants (10, 11). It has multiple pharmacological activities, including antibacterial (12), antitumor, antiulcer (13), myocardial-protecting (14), vasoprotective, immunomodulatory, antioxidant, cytoprotective, and neuroprotective activities (15). Moreover, in the reproductive system, rutin has shown a possible protection of the testicular tissue and reproduction from oxidative stress in diabetes mellitus (16) and has led to the amelioration of cisplatin-induced reproductive toxicity (17). Considerable interest has recently been directed to the role and usage of natural antioxidants as a means of preventing oxidative damage in different conditions with a high oxidative stress (13).

Steroids play a prominent role in the acceptance of transplanted tissues. Extensive evidence suggests that estrogen directly modulates angiogenesis via effects on endothelial cells (18). Estrogen,  $17\beta$ -estradiol ( $E_2$ ), plays an important role in regulating the proliferation and maturation of ovarian follicles. It has a critical function in the female reproductive system (19, 20). Its effects are mediated through two types of receptor: Estrogen Receptor alpha ( $ER\alpha$ ) and Estrogen Receptor beta ( $ER\beta$ ). These nuclear receptors mediate the biological effects of estrogens and anti-estrogens and act mainly in the regulation of the Estrogen Receptor (ER) target-gene expression (21). Progesterone ( $P_4$ ) is an endogenous steroid involved in the menstrual cycle and pregnancy (22). The physiological effects of  $P_4$  are exerted by the progesterone receptor (PR), a member of the nuclear receptor superfamily of transcription factors (23).

## 2. Objectives

The above-mentioned facts form the basis for studying whether the antioxidant mechanisms involved in rutin-

mediated protection from ischemic damage after ovarian transplantation can improve the conditions of fertility. Consequently, the present animal experimental study aimed to investigate the effects of different concentrations of rutin on the return of fertility after ovarian heterotopic allograft and the expression of  $ER\alpha$ ,  $ER\beta$ , and PR in Balb/c mice.

## 3. Methods

### 3.1. Animals

The present animal experimental study was conducted on 40 adult female Balb/c mice, 5 - 6 weeks of age, weighing approximately 25 - 35 g. Animals were obtained from the Animal House of Jundishapur University of Medical Sciences, Ahvaz, Iran, and maintained at standard conditions (temperature 22 - 24°C and humidity 55% - 65% in a 12-hour light-dark cycle) during 2016 - 2018 in Jundishapur University of Medical Sciences, Ahvaz, Iran. Animals had free access to sufficient amounts of water and food during the study. This study was performed in accordance with the guidelines for Animal Research of the National Institutes of Health (NIH, Bethesda, MD, USA). The ethical code was approved by the Ethics Committee of Jundishapur University of Medical Sciences, Ahvaz, Iran, on 16 May, 2017 (Code: IR.AJUMS.REC.1396.40). For mating, a male and a female animal were housed in plastic cages. Ten days after birth, the newborn female mice were selected for the donation of the ovary to adult female mice. According to a similar study by Domitrovic et al. (24) in 2012, the mean relative weight of the liver in the control group was  $5.0 \pm 7.4$  and, in the intervention group with  $CCl_4$ , it was  $4.0 \pm 7.6$ . By considering the confidence level of 95% and a test power of 80%, five mice for each study group were estimated using the following formula.

$$n = \frac{(s_1^2 + s_2^2) \left( z_{1-\frac{\alpha}{2}} + z_{1-\beta} \right)^2}{(\bar{x}_1 - \bar{x}_2)^2}$$

The animals were divided into eight groups by permuted block randomization. The eight groups consisted of OVX (ovariectomy/negative control), undergoing ovariectomy; the normal group (positive control), normal and untreated mice; OVX + OVA (ovariectomy and transplantation) (control), undergoing ovariectomy and transplantation, treated with 0.5 mL of normal saline; OVX + OVA + Rut 10, OVX + OVA + Rut 30, OVX + OVA + Rut 60, and OVX + OVA + Rut 100, undergoing ovariectomy and transplantation, treated with 10, 30, 60, and 100 mg/kg of rutin, respectively; and autograft, without any treatment. All groups were treated once a day for two weeks with the intraperitoneal injection of normal saline for the control group

and rutin for other groups. Rutin was purchased from the Sigma Aldrich Company (St. Louis, MO) and dissolved in DMSO. All mice were anesthetized using ketamine (60 mg/kg) and xylazine (6 mg/kg) and then ovariectomized in sterile conditions. One week after ovariectomy, two ovaries were transplanted under the muscles behind the neck.

### 3.2. Sample Collection

After transplantation, to ensure that the estrus phase is resumed in the OVX + OVA mice, the vaginal smear was prepared every day, 3 - 4 days after transplantation. For serum collection, 24h after the last dose of rutin, 1.5 mL of blood was collected from the heart. The blood sample was centrifuged at 3000 g for 5 min, and the serum was separated. Then, all samples were kept at -20 °C until use for the hormonal assay.

### 3.3. Hormonal Assay

Serum E<sub>2</sub> and P<sub>4</sub> levels were analyzed by the ELISA method in duplicate using a progesterone kit (Crystal Chem, USA) with the sensitivity of 0.04 ng/mL and assay range of 0.4 - 100 ng/mL; and an estrogen kit (Cusabio, USA) with the sensitivity of 40 pg/mL and assay range of 40 - 1500 pg/mL, according to the manufacturer's instructions.

### 3.4. Histology

Two weeks after transplantation, all grafted ovaries were collected and fixed in 10% formalin and then the tissues were embedded in paraffin, serially sectioned at 5 μm, and stained with the Haematoxylin and Eosin. Ovaries of age-matched 32-day-old mice were histologically processed as described above as the in-vivo development of normal control. Microphotography slides were obtained using a light microscope (Olympus CH-BI45-2) with the magnification of 20X and 40X and assessed by three experts. Primordial, primary, preantral, antral and atretic follicles were counted.

### 3.5. qRT-PCR

Total RNA was extracted from endometrial tissue using RiboEx (GeneAll Biotechnology Co, USA) according to the protocol of the manufacturer. Extracted RNA was kept at -80°C. For cDNA synthesis, 500 ng of total RNA was added to the cDNA synthesis tube (Thermo Scientific, USA). The program of cDNA synthesis was: 30°C for 5 min, 42°C for 1 h, 70°C for 5 min, heated at 95°C for 5 min, and stored at -20°C.

qRT-PCR primers were designed using AlleleID 6.0 software and ordered to the CinnaGen Company (Tehran, Iran) for commercial synthesis. The primer sequences of the gene of interest are fully described in Table 1. qRT-PCR reactions were performed in the ABI StepOne instrument (Applied Biosystems, USA) under the following conditions:

**Table 1.** List of the Primer Sequences of Interest Gene

Gene	Primers
<b>GAPDH</b>	
FW	5'-AGCAAGGACACTGAGCAAGAG-3'
RE	5'-GGATGGAAATTGTGAGGGAGATG-3'
<b>ER<math>\alpha</math></b>	
FW	5'-TAGCGGCAACAGTGAATCC-3'
RE	5'-TGGCAAGGTAAGCAATGGC-3'
<b>ER<math>\beta</math></b>	
FW	5'-ATGGACTGTAGAACGGTGTGG-3'
RE	5'-GTGAGGTAGGAATGCCAAACG-3'
<b>PR</b>	
FW	5'-GATTCAGAAGCCAGCCAGAG-3'
RE	5'-CACAGGTAAGCACGCCATAG-3'

95°C for 15 s and 60°C for 1 min for up to 40 cycles. The data are presented as the relative quantity of normalized-target RNA using the  $\Delta\Delta$ CT method (25) (internal control gene GAPDH). Each sample was examined in duplicate.

### 3.6. Western Blotting

The expression of ER $\alpha$ , ER $\beta$ , and PR was assessed by the Western blot technique. It was prepared using the lysis buffer containing the RIPA buffer and a protease inhibitor cocktail. Total protein was measured by the Bradford method. Equal amounts of protein were resolved by 12% SDS-polyacrylamide gel electrophoresis using the Bio-Rad system (Mini-PROTEAN Tetra System, USA), and then transferred on the PVDF membrane (Immobilon-p transfer membranes) using the wet transfer method. The membrane was blocked overnight with casein 1% in Tris Buffered Saline (TBS) with Tween-20. The membrane was probed with primary antibody (Abcam, USA) and then with horseradish peroxidase-conjugated secondary antibody (Abcam, USA). Finally, ECL plus was used for the final detection of protein bands, and all data were analyzed using the ImageJ software.

### 3.7. Statistical Analysis

The data were analyzed using IBM SPSS Statistic Software for Windows, version 22.0 (IBM Corp., Armonk, N.Y., USA). All devices and equipment were calibrated to obtain unbiased data during the study. In this study, descriptive statistics, including mean, standard deviation, and error bar were used to plot the mean and standard deviation of each group. Descriptive data are presented as a mean  $\pm$  standard deviation. The normality of variables was examined using the Shapiro-Wilk test (P value > 0.05). Since the data showed a normal distribution, one-way ANOVA with

a post-hoc test was used. In this study, to determine the agreement between the two observers, the kappa coefficient was calculated which was 0.80, indicating that the agreement rate is reasonable. The P value < 0.05 was considered statistically significant.

## 4. Results

### 4.1. Level of E<sub>2</sub> and P<sub>4</sub>

As illustrated in Figure 1, the serum level of E<sub>2</sub> and P<sub>4</sub> dramatically decreased in the OVX group in comparison with the control group. In OVX, OVX + OVA, OVX + OVA Rut 10, OVX + OVA Rut 30, OVX + OVA Rut 60, OVX + OVA Rut 100, autograft and normal groups, the level of E<sub>2</sub> was, 0, 25, 32.6, 31, 21.2, 23.3, 28, 26.3 (pg/mL) respectively and the level of P<sub>4</sub> was 0.4, 0.4, 2.9, 2.93, 1.26, 6.13, 6.5, 8 (ng/mL), respectively. The level of E<sub>2</sub> increased in all OVA + rutin groups and the autograft group in comparison with the OVX group (P value < 0.01). Intra-group analysis of the relation between various concentrations of rutin showed the most increase in 0, 10, 30, 100, and the autograft groups. As expected, the level of P<sub>4</sub> was increased in all OVA + rutin groups, in addition to the autograft group, compared to the OVX + OVA group. Intra-group analysis indicated the less effect of OVX + rutin 60 compared to OVX + rutin 0, 10, 30, 100, and also the autograft group (P value < 0.0001) (Table 2).

Table 2. Changes in Sex Hormones in Groups Receiving Rutin<sup>a</sup>

Groups	Estrogen, pg/mL	Progesteron, ng/mL
OVX	0.0 ± 0.0 <sup>b</sup>	0.4667 ± 0.1155
Normal	26.33 ± 4.726	8.000 ± 1.600 <sup>b</sup>
Treat 0	25.00 ± 10.15	0.4667 ± 0.2517
Treat 10	32.67 ± 4.163	2.900 ± 0.5196
Treat 30	31.00 ± 2.000	2.933 ± 0.6429
Treat 60	21.23 ± 14.03	1.267 ± 0.6658
Treat 100	23.33 ± 0.5774	6.133 ± 1.026 <sup>b</sup>
Autograft	28.00 ± 6.245	6.500 ± 2.166 <sup>b</sup>

<sup>a</sup>Values are expressed as mean ± SD.

<sup>b</sup>Significant difference compared to treat 0 (P < 0.05).

### 4.2. Follicle Development

The presence of numerous cysts in all groups under treatment with rutin was observed. The number of these cysts was decreased with a decrease in the dose of rutin. The number of primordial follicles in treated groups was in the following order: treat 60 > treat 0 > treat 30 > treat 10 > treat 100. Moreover, the number of primary follicles in treated groups was in the following order: treat 60 > treat 10 > treat 30 > treat 100 > treat 0. Also, the number of pre-antral follicles in the treated groups was in the following

order: treat 10 > treat 60 > treat 30 > treat 0 > treat 100 (P value < 0.0001). In addition, the number of antral follicles in the treated groups was in the following order: treat 10 > treat 30 > treat 60 > treat 0 > treat 100 (P value < 0.0001). Finally, the number of atretic follicles in the treated groups was in the following order: treat 60 > treat 100 > treat 30 > treat 10 = treat 0 (P value < 0.001) (Figure 2 and Table 3).

### 4.3. PR, ER $\alpha$ , and ER $\beta$ mRNA Expression

Results indicated that, as expected, the PR, ER $\alpha$ , and ER $\beta$  mRNA expression in the OVX group was almost zero. OVX + rutin 10 and 30 groups had the highest levels of ERs mRNA (P value < 0.001). The mean mRNA expression level of ER $\alpha$  was 0.3513 ± 0.5687, 1.000 ± 0.0, 5.430 ± 1.000, 5.218 ± 1.000, 2.250 ± 1.000, 2.143 ± 1.100, 5.429 ± 1.000 and 5.760 ± 1.000 (Figure 3A); The mean mRNA expression level of ER $\beta$  was 0.3344 ± 0.1000, 1.000 ± 0.0, 3.608 ± 0.5774, 4.497 ± 1.000, 2.877 ± 1.528, 3.442 ± 1.000, 0.4454 ± 0.5774 and 2.998 ± 1.000 (Figure 3B) and The mean mRNA expression level of PR was 0.6776 ± 0.5774, 1.000 ± 0.0, 1.073 ± 1.000, 1.101 ± 1.000, 1.733 ± 1.000, 1.018 ± 1.000, 1.038 ± 1.000 and 3.528 ± 1.000 in OVX, OVX + OVA, OVX + OVA Rut 10, OVX + OVA Rut 30, OVX + OVA Rut 60, OVX + OVA Rut 100, autograft and normal, respectively. In all the treated groups, an increase in PR mRNA versus the OVX group was shown, but there was no statistically significant difference (Figure 3C).

### 4.4. Protein Expression Level of PR, ER $\alpha$ , and ER $\beta$

The protein expression level of PR, ER $\alpha$ , and ER $\beta$  within the endometrial tissue of the mice had the same pattern across groups. In OVX, OVX + OVA, OVX + OVA Rut 10, OVX + OVA Rut 30, OVX + OVA Rut 60, OVX + OVA Rut 100, autograft and normal, the level of ER $\alpha$  was, 0.0044, 0.81, 1.88, 1.78, 1.46, 0.56, 1.76, 1.65, respectively. The level of ER $\beta$  was 0.025, 0.14, 1.25, 1.05, 0.56, 0.80, 0.19, 0.50, respectively, and the level of PR was 0.09, 0.21, 0.14, 0.88, 0.10, 0.28, 0.09, and 0.59 respectively. The rates in the OVX + rutin 100 and 0 groups had the lowest levels of ERs (P value < 0.05). The highest level of ER $\alpha$  and ER $\beta$  was seen in OVX + rutin 10 and 30. There were no detectable expressions in the OVX group (P value < 0.01). Moreover, the protein expression of PR in all OVX + OVA groups treated with rutin was increased in comparison with the OVX group, but there was no significant difference (Figures 3D - 3G and Table 4).

## 5. Discussion

Results of the present study demonstrate that fresh hetero allograft ovarian transplantation with different concentrations of rutin in mice can restore ovarian function, as evidenced by the production of E<sub>2</sub> and P<sub>4</sub> along




Figure 1. Serum ELISA results for estrogen and progesterone

Table 3. Follicular Density in Grafted Ovaries<sup>a</sup>

Groups	No. of Follicles				
	Primordial	Primary	Preantral	Antral	Atretic
Normal	19.67 ± 5.686	19.67 ± 5.686	21.00 ± 1.000 <sup>b</sup>	23.00 ± 1.000 <sup>b</sup>	5.000 ± 1.000 <sup>b</sup>
Treat 0	17.67 ± 2.517	16.67 ± 2.082	0.3333 ± 0.5774	0.3333 ± 0.5774	12.33 ± 2.517
Treat 10	12.33 ± 2.517	17.67 ± 2.517	13.00 ± 1.000 <sup>b</sup>	10.67 ± 3.055 <sup>b</sup>	12.33 ± 2.517
Treat 30	13.67 ± 3.786	17.33 ± 2.082	1.000 ± 1.000	1.667 ± 1.528	19.00 ± 1.000 <sup>b</sup>
Treat 60	19.67 ± 1.528	19.67 ± 1.528	3.000 ± 1.000 <sup>b</sup>	1.000 ± 1.000	25.00 ± 5.000 <sup>b</sup>
Treat 100	12.00 ± 3.000	17.00 ± 4.359	0.0 ± 0.0	0.0 ± 0.0	23.00 ± 2.646 <sup>b</sup>

<sup>a</sup>Values are expressed as mean ± SD.

<sup>b</sup>Significant difference compared to treat 0 ( $P < 0.05$ ).

Table 4. Protein Expression Level of Estrogen Receptor Alpha (ER $\alpha$ ), Estrogen Receptor Beta (ER $\beta$ ), and Progesterone Receptor (PR)<sup>a</sup>


Groups	ERa	ERb	PR
OVX	0.007864 ± 0.003837	0.05174 ± 0.03778	0.1156 ± 0.06584
Normal	1.495 ± 0.3662 <sup>b</sup>	0.6670 ± 0.1752 <sup>b</sup>	0.6010 ± 0.1626 <sup>b</sup>
Treat 0	0.7089 ± 0.1131	0.1534 ± 0.06109	0.1423 ± 0.09290
Treat 10	1.615 ± 0.1701 <sup>b</sup>	0.7747 ± 0.4365 <sup>b</sup>	0.1593 ± 0.06674
Treat 30	1.744 ± 0.1779 <sup>b</sup>	0.9220 ± 0.1245 <sup>b</sup>	0.2064 ± 0.1529
Treat 60	0.9776 ± 0.7617	0.7701 ± 0.2150 <sup>b</sup>	0.1591 ± 0.06259
Treat 100	0.4953 ± 0.2098	0.6676 ± 0.1547 <sup>b</sup>	0.2763 ± 0.05559
Autograft	1.770 ± 0.1000 <sup>b</sup>	0.1860 ± 0.02788	0.2537 ± 0.1264

<sup>a</sup>Values are expressed as mean ± SD.

<sup>b</sup>Significant difference compared to treat 0 ( $P < 0.05$ ).

with the expression of ER $\alpha$ , ER $\beta$ , and PR in the transplanted mice. Also, results showed that the use of rutin in allograft for restoring the ovarian function could have an equal value to autograft transplantation in some concentrations.

Beazley and Nurminskaya explored the effects of flavonoid on fertility in female mice. They demonstrated that quercetin leads to a 60% reduction in the number of litters, but enhances folliculogenesis in the ovaries of female offspring (26). In this study, consistent with the study



**Figure 2.** Number of follicles in examining the ovaries in normal, OVX + OVA, OVX + OVA + Rut 10, OVX + OVA + Rut 30, OVX + OVA + Rut 60, and OVX + OVA + Rut 100 groups; (A) number of primordial follicles, (B) number of primary follicles, (C) number of preantral follicles, (D) number of antral follicles, and (E) number of atretic follicles.


by Beazley and Nurminskaya, rutin as a flavonoid caused an increase in the maturation of follicles but was not able to increase fertility (increased atretic follicles).

Treatment of OVX + OVA mice with rutin possibly restored the estrous cyclicity. However, the rutin-treated groups at different concentrations showed different serum levels of  $E_2$  and  $P_4$ .

In the present study,  $E_2$  concentration in plasma was

significantly increased. Guo et al. in a similar study reported that the administration of rutin has an effect on ovariectomized rats similar to the administration of  $E_2$ . They concluded that rutin is able to increase  $E_2$  concentration in serum and mammary glands (27). In another study on flavonoids, Tan et al. demonstrated the same role of these molecules in increasing  $E_2$  levels in pubertal female rats (28). In addition, Nynca et al. reported genistein action





**Figure 3.** mRNA expression level of ER $\alpha$ , ER $\beta$ , and PR (A, B, C); protein expression level of ER $\alpha$ , ER $\beta$ , and PR (D, E, F, G)

(a flavonoid) on E<sub>2</sub> production by porcine granulosa cells of medium follicles. All doses of genistein increased basal E<sub>2</sub> secretion by the granulosa cells cultured for 48 h (29). In contrast, daidzein as another flavonoid did not alter the granulosa secretion of E<sub>2</sub> (29).

P<sub>4</sub> was also studied as a hormone with fluctuations in

ovary activity and follicle growth. Interestingly, the level of P<sub>4</sub> was higher in the autograft group than rutin10, 30, and 60 groups, while there was no significant difference in comparison with rutin 100. Jahan et al. in a study on rats with polycystic ovary syndrome demonstrated an increase in P<sub>4</sub> due to the administration of rutin at two doses

of 100 mg/kg and 150 mg/kg (30). In contrast, in other studies, genistein inhibited P<sub>4</sub> production by the porcine granulosa cells of medium follicles (29) and porcine luteinized granulosa cells isolated from large follicles (31).

Results showed that ER $\beta$  expression in the endometrial tissue was significantly increased in all OVX + OVA groups treated with rutin versus OVX and treat O groups. As demonstrated in the E<sub>2</sub> pattern, an increase in ER $\beta$  in the treated group had a significantly higher protein expression level than the autograft group, suggesting that rutin increased ER $\alpha$  expression in the endometrial tissue. The protein expression level of ER $\alpha$  has the same pattern as ER $\beta$  in all treated groups in comparison with control, OVX, and OVX + OVA groups. A prominent difference was found in the expression level of ER $\alpha$  which was considerably higher in the autograft group in comparison with all other groups. In their study on the possible pathway using which rutin can ameliorate oxidative injury, Hong et al. report that rutin can attenuate the ischemia/reperfusion injury in ovariectomized rats via ER-mediated signaling pathways (BDNF-TrkB and NGF-TrkA signaling pathway) (32). The structural similarity of rutin to endogenous estrogen and plant estrogen makes this molecule able to be absorbed by target cells, be bound to ER, and then exert estrogen-like effects (33). In addition to the direct effects of rutin on ER activation and signaling initiation, this agent is able to increase the production of estrogen directly, as mentioned by Guo (27). Similarly, Nynca et al. demonstrated that genistein caused a significant increase in the ER $\beta$  mRNA level in granulosa cells of large follicles (31) and granulosa cells of medium follicles (29). However, it did not alter ER $\alpha$  mRNA level in the culture porcine granulosa cells (29, 31). In contrast, the expression of ER $\beta$  protein was affected by genistein in the granulosa cells of medium follicles (29), but it was not detected in the granulosa cells of large follicles (31). Daidzein decreased mRNA expression of ER $\alpha$  in medium follicles, but the expression of ER $\beta$  mRNA was not affected by daidzein. ER $\alpha$  protein was not detected while ER $\beta$  protein was found in the nuclei of the cells. Daidzein upregulated the expression of ER $\beta$  protein in the cells (29).

The PR expression in the endometrial tissue had a decreased pattern in comparison with the control group. The maximum protein expression level of PR was observed in the treat rutin 100 group. In all transplanted groups, an increase of PR versus the OVX group was shown. However, in groups 0, 10, 30, 60, and the autograft group, there was no statistically significant difference. These results suggest that rutin is able to enhance PR. Parallel with this study; Rosenberg et al. demonstrated that flavonoids have a progesterone-like antagonist activity. In their study on PR in breast cancer cell lines, it was shown that flavonoids bind to the PR and act as a blocker. They demonstrated that flavonoids could modulate PR expression (34). It seems

that rutin, as a flavonoid component, has a modulatory effect on PR beside its antioxidant and estrogen-like abilities. In agreement with Rosenberg, no statistically significant change in PR was observed in the present study.

## 5.1. Strengths and Limitations

### 5.1.1. Limitations

- The sample size was small (n = 5) for each group.
- No additional evaluation methods such as immunohistochemistry were used.

### 5.1.2. Strengths

- Evaluation methods such as qRT-PCR and Western blotting were employed.
- Four dosages of rutin were tested.

## 5.2. Conclusions

The present study demonstrates that the effects of rutin on restoring the estrous cyclicity after transplantation rely on its antioxidant effect on the inhibition of the increased oxidative stress. The results of the present study indicated that rutin increased the E<sub>2</sub> and P<sub>4</sub> levels in ovarian hetero allograft mice. Rutin also upregulated the expression of ER $\alpha$  and ER $\beta$  but had no significant effect on PR. ER upregulation led to an enhanced function of estrogen, improved the engraftment and function of the transplanted ovarian tissue, and restored estrous cyclicity.

## Footnotes

**Authors' Contribution:** Ghasem Saki supervised the research. Aligholi Sobhani supervised the research and edited. Masoud Hemadi designed the project. Tayebeh Hadigol obtained the specimens, carried out the experiment, and writing the manuscript. Saeid Nekoonam analyzed the data. Maryam Dastoorpoor analyzed the statistics section. Alireza Shams carried out the experiment. Bahram Eslami Farsani writing the manuscript.

**Conflict of Interests:** The authors declare no conflict of interests.

**Ethical Approval:** IR.AJUMS.REC.1396.40.

**Funding/Support:** This article and thesis supported by Ahvaz Jundishapur University of Medical Sciences.

## References

1. Sheikhi M, Hulthenby K, Niklasson B, Lundqvist M, Hovatta O. Clinical grade vitrification of human ovarian tissue: An ultrastructural analysis of follicles and stroma in vitrified tissue. *Hum Reprod.* 2011;26(3):594-603. doi: 10.1093/humrep/deq357. [PubMed: 2121741]. [PubMed Central: PMC3037791].



2. Callejo J, Vilaseca S, Ordi J, Cabre S, Laila JM, Balasch J. Heterotopic ovarian transplantation without vascular pedicle in syngeneic Lewis rats: Long-term evaluation of effects on ovarian structure and function. *Fertil Steril*. 2002;77(2):396-402. [PubMed: [11821104](#)].
3. Nayki C, Nayki U, Keskin Cimen F, Kulhan M, Yapca OE, Kurt N, et al. The effect of rutin on ovarian ischemia-reperfusion injury in a rat model. *Gynecol Endocrinol*. 2018;34(9):809-14. doi: [10.1080/09513590.2018.1450378](#). [PubMed: [29564969](#)].
4. Yaman Tunc S, Agacayak E, Goruk NY, Icen MS, Turgut A, Alabalik U, et al. Protective effects of honokiol on ischemia/reperfusion injury of rat ovary: An experimental study. *Drug Des Devel Ther*. 2016;10:1077-83. doi: [10.2147/DDDT.S93768](#). [PubMed: [27022246](#)]. [PubMed Central: [PMC4789835](#)].
5. Esteban-Zubero E, Garcia-Gil FA, Lopez-Pingarron L, Alatorre-Jimenez MA, Inigo-Gil P, Tan DX, et al. Potential benefits of melatonin in organ transplantation: A review. *J Endocrinol*. 2016;229(3):R129-46. doi: [10.1530/JOE-16-0117](#). [PubMed: [27068700](#)].
6. Lawson M, Jomova K, Poprac P, Kuca K, Musilek K, Valko M. Nutritional antioxidant therapies: Treatments and perspective. *Free radicals and antioxidants in human disease*. Springer; 2017. p. 283-305. doi: [10.1007/978-3-319-67625-8\\_12](#).
7. Kawamura K, Cheng Y, Sun YP, Zhai J, Diaz-Garcia C, Simon C, et al. Ovary transplantation: To activate or not to activate. *Hum Reprod*. 2015;30(11):2457-60. doi: [10.1093/humrep/dev211](#). [PubMed: [26345683](#)].
8. Silber S. Ovarian tissue cryopreservation and transplantation: Scientific implications. *J Assist Reprod Genet*. 2016;33(12):1595-603. doi: [10.1007/s10815-016-0814-1](#). [PubMed: [27722934](#)]. [PubMed Central: [PMC5171890](#)].
9. Gavish Z, Spector I, Peer G, Schlatt S, Wistuba J, Roness H, et al. Follicle activation is a significant and immediate cause of follicle loss after ovarian tissue transplantation. *J Assist Reprod Genet*. 2018;35(1):61-9. doi: [10.1007/s10815-017-1079-z](#). [PubMed: [29098533](#)]. [PubMed Central: [PMC5758475](#)].
10. Agustin-Salazar S, Medina-Juarez LA, Soto-Valdez H, Manzaneres-Lopez F, Gamez-Meza N. Influence of the solvent system on the composition of phenolic substances and antioxidant capacity of extracts of grape (*Vitis vinifera* L.) marc. *Aust J Grape Wine Res*. 2014;20(2):208-13. doi: [10.1111/ajgw.12063](#).
11. Cruz-Zuniga JM, Soto-Valdez H, Peralta E, Mendoza-Wilson AM, Robles-Burgueno MR, Auras R, et al. Development of an antioxidant biomaterial by promoting the deglycosylation of rutin to isoquercetin and quercetin. *Food Chem*. 2016;204:420-6. doi: [10.1016/j.foodchem.2016.02.130](#). [PubMed: [26988520](#)].
12. Ananth DA, Rameshkumar A, Jeyadevi R, Jagadeeswari S, Nagarajan N, Renganathan R, et al. Antibacterial potential of rutin conjugated with thioglycolic acid capped cadmium telluride quantum dots (TGA-CdTe QDs). *Spectrochim Acta A Mol Biomol Spectrosc*. 2015;138:684-92. doi: [10.1016/j.saa.2014.11.082](#). [PubMed: [25544184](#)].
13. Ganeshpurkar A, Saluja AK. The pharmacological potential of rutin. *Saudi Pharm J*. 2017;25(2):149-64. doi: [10.1016/j.jsps.2016.04.025](#). [PubMed: [28344465](#)]. [PubMed Central: [PMC5355559](#)].
14. Lin Q, Chen XY, Zhang J, Yuan YL, Zhao W, Wei B. Upregulation of SIRT1 contributes to the cardioprotective effect of rutin against myocardial ischemia-reperfusion injury in rats. *J Funct Food*. 2018;46:227-36. doi: [10.1016/j.jff.2018.05.007](#).
15. Almutairi MM, Alanazi WA, Alshammari MA, Alotaibi MR, Alhoshani AR, Al-Rejaie SS, et al. Neuro-protective effect of rutin against Cisplatin-induced neurotoxic rat model. *BMC Complement Altern Med*. 2017;17(1):472. doi: [10.1186/s12906-017-4976-9](#). [PubMed: [28962559](#)]. [PubMed Central: [PMC5622464](#)].
16. Butchi Akondi R, Kumar P, Annapurna A, Pujari M. Protective effect of rutin and naringin on sperm quality in streptozotocin (STZ) induced type 1 diabetic rats. *Iran J Pharm Res*. 2011;10(3):585-96. [PubMed: [24250392](#)]. [PubMed Central: [PMC3813033](#)].
17. Aksu EH, Kandemir FM, Ozkaraca M, Omur AD, Kucukler S, Comakli S. Rutin ameliorates cisplatin-induced reproductive damage via suppression of oxidative stress and apoptosis in adult male rats. *Andrologia*. 2017;49(1). doi: [10.1111/and.12593](#). [PubMed: [27106863](#)].
18. Losordo DW, Isner JM. Estrogen and angiogenesis: A review. *Arterioscler Thromb Vasc Biol*. 2001;21(1):6-12. [PubMed: [11145928](#)].
19. Xu Y, Ding J, Ma XP, Ma YH, Liu ZQ, Lin N. Treatment with panax ginseng antagonizes the estrogen decline in ovariectomized mice. *Int J Mol Sci*. 2014;15(5):7827-40. doi: [10.3390/ijms15057827](#). [PubMed: [24806341](#)]. [PubMed Central: [PMC4057706](#)].
20. Kim MH, Kim HM, Jeong HJ. Estrogen-like osteoprotective effects of glycine in in vitro and in vivo models of menopause. *Amino Acids*. 2016;48(3):791-800. doi: [10.1007/s00726-015-2127-6](#). [PubMed: [26563333](#)].
21. Debeb YG, Berihu BA. Review on the role of estrogen receptors in breast cancer. *Int J Pharm Sci Res*. 2015;6(8):1100-3.
22. Jeon SY, Hwang KA, Choi KC. Effect of steroid hormones, estrogen and progesterone, on epithelial mesenchymal transition in ovarian cancer development. *J Steroid Biochem Mol Biol*. 2016;158:1-8. doi: [10.1016/j.jsbmb.2016.02.005](#). [PubMed: [26873134](#)].
23. Wu SP, Li R, DeMayo FJ. Progesterone receptor regulation of uterine adaptation for pregnancy. *Trends Endocrinol Metab*. 2018;29(7):481-91. doi: [10.1016/j.tem.2018.04.001](#). [PubMed: [29705365](#)]. [PubMed Central: [PMC6004243](#)].
24. Domitrovic R, Jakovac H, Vasiljev Marchesi V, Vladimir-Knezevic S, Cvijanovic O, Tadic Z, et al. Differential hepatoprotective mechanisms of rutin and quercetin in CCl(4)-intoxicated BALB/cN mice. *Acta Pharmacol Sin*. 2012;33(10):1260-70. doi: [10.1038/aps.2012.62](#). [PubMed: [22902988](#)]. [PubMed Central: [PMC4002705](#)].
25. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc*. 2008;3(6):1101-8. [PubMed: [18546601](#)].
26. Beazley KE, Nurminskaya M. Effects of dietary quercetin on female fertility in mice: Implication of transglutaminase 2. *Reprod Fertil Dev*. 2016;28(7):974-81. doi: [10.1071/RD14155](#). [PubMed: [25557047](#)]. [PubMed Central: [PMC4794435](#)].
27. Guo XD, Diao QY, Wang YY, Tu Y, Deng KD, Wang XJ, et al. The effect of administration of rutin on plasma levels of estrogen, prolactin, growth hormone and gene expression of their receptors in mammary glands in ovariectomized rats. *J Integr Agr*. 2012;11(10):1700-6. doi: [10.1016/s2095-3119\(12\)60173-7](#).
28. Tan KP, Chen J, Ward WE, Thompson LU. Mammary gland morphogenesis is enhanced by exposure to flaxseed or its major lignan during suckling in rats. *Exp Biol Med (Maywood)*. 2004;229(2):147-57. [PubMed: [14734793](#)].
29. Nynca A, Slonina D, Jablonska O, Kaminska B, Ciereszko RE. Daidzein affects steroidogenesis and oestrogen receptor expression in medium ovarian follicles of pigs. *Acta Vet Hung*. 2013;61(1):85-98. doi: [10.1556/AVet.2012.060](#). [PubMed: [23439294](#)].
30. Jahan S, Munir F, Razak S, Mehboob A, Ain QU, Ullah H, et al. Ameliorative effects of rutin against metabolic, biochemical and hormonal disturbances in polycystic ovary syndrome in rats. *J Ovarian Res*. 2016;9(1):86. doi: [10.1186/s13048-016-0295-y](#). [PubMed: [27923406](#)]. [PubMed Central: [PMC5142269](#)].
31. Nynca A, Sadowska A, Orłowska K, Jablonska M, Ciereszko RE. The effects of phytoestrogen genistein on steroidogenesis and estrogen receptor expression in porcine granulosa cells of large follicles. *Folia Biol (Krakow)*. 2015;63(2):119-28. [PubMed: [26255463](#)].
32. Liu H, Zhong L, Zhang Y, Liu X, Li J. Rutin attenuates cerebral ischemia-reperfusion injury in ovariectomized rats via estrogen-receptor-mediated BDNF-TrkB and NGF-TrkA signaling. *Biochem Cell Biol*. 2018;96(5):672-81. doi: [10.1139/bcb-2017-0209](#). [PubMed: [29420916](#)].
33. Artursson P, Palm K, Luthman K. Caco-2 monolayers in experimental and theoretical predictions of drug transport. *Adv Drug Deliv Rev*. 2001;46(1-3):27-43. [PubMed: [11259831](#)].
34. Rosenberg RS, Grass L, Jenkins DJ, Kendall CW, Diamandis EP. Modulation of androgen and progesterone receptors by phytochemicals in breast cancer cell lines. *Biochem Biophys Res Commun*. 1998;248(3):935-9. doi: [10.1006/bbrc.1998.8977](#). [PubMed: [9704030](#)].